Notes and Observations

A FURTHER RECORD OF DORYTOMUS SALICINUS (GYLLENHAL) (COL.: CURCULIONIDAE) FROM CUMBRIA. — Since recording Dorytomus salicinus (Gyllenhal) for the first time from Cumbria and vice county 70 Cumberland (1979, Entomologist's Record, 91: 27-28). I have recently discovered this rare and very local weevil in a second locality in the county. While collecting phytophagous coleoptera on Braithwaite Moss, 10km square NY22 at the North east end of Bassenthwaite Lake, Cumbria on April 4th 1981, I beat four specimens of D. salicinus from two somewhat isolated trees of Salix atrocinerea Brot. growing in an open area of marshy ground and near to the edge of a small conifer wood. The Salix trees contained many catkins and a good number of developing leaf shoots were also present, and it is possible that the weevils were feeding on these structures and perhaps oviposition was also taking place at this time in the flowers.

I also took by beating from the above trees of sallow a few specimens of the rather common *Dorytomus taeniatus* (Fabricius), and I observed that while the two species were in the net that *D. salicinus* became more active only after a very brief feining period, while *D. taeniatus* remained motionless and quiescent for a comparatively longer period and it was generally observed to

walk slower.

Among other beetles collected at this site were two specimens of *Dorytomus melanophthalmus* (Paykull) also beaten from sallow, and one *Anthonomus bituberculatus* Thomson, C. G. I also took the somewhat local chrysomelid *Aphthona nonstriata* (Goeze) on yellow iris. — R. W. J. READ, 43 Holly Terrace, Hensingham, Whitehaven, Cumbria, CA28 8RF.

SAMUEL DALE'S "1704" CATALOGUE OF ENGLISH BUTTER-FLIES. – John Ray and James Petiver have at least partially received their due as founders of the scientific study of insects, especially Lepidoptera, in England. Their colleague Samuel Dale (ca. 1659-1739), a Braintree, Essex apothecary and close friend and neighbour of Ray, has had relatively little notice. Dale did not attain the status of Ray or Petiver, but he contributed to both of their collections as an assiduous field entomologist, and was one of the several instigators of the eventual publication of Ray's great post-

humous work, Historia Insectorum (London, 1710).

In the library of the Royal Entomological Society of London is Dale's copy of Martin Lister's extremely rare English edition of Johannes Godartius, Of Insects (York, 1682; Lisney 45; only 150 copies were printed), also ex libris James Francis Stephens. Bound before the printed work is a manuscript in Dale's hand, "A Cataloge [sic] of English Butterflies Reduced to Mr. Rays Method 1704," but containing later additions. The butterflies are described in short Latin sentences with references to published sources. The manuscript includes Dale's entries concerning other insects, and one note is dated as late as 1728. Numerous Rhopalo-

cera are recognisable from Dale's descriptions, and other characterisations are now being studied so that an analysis of the manuscript can be included in a forthcoming contribution on the role of Samuel Dale in the history of British entomology. — R. S. WILKINSON, The American Museum of Natural History, New York, New York

10024.

AGONOPTERIX ASTRANTIAE (HEINEMANN) IN NORTH WALES IN 1981 - Further to my paper on Oecophoridae (Ent. Rec., 93:60) and that of Heckford and Langmaid. (Ent. Rec. 93, 100), I found three larvae on Sanicula on 22nd June in mixed deciduous woods on carboniferous limestone near Llandudno (VC 49). Two were almost full fed and spun up in the tissue paper provided within five days; the third larva also well grown produced a parasite. All larvae were in the spun edge of a leaf rolled upwards as described by Heckford and Langmaid and there were signs of spinning and feeding on nearby leaves on the same or adjacent plants. In the breeding box, while the larvae remained in their original folds in daytime, other leaves were eaten and frass scattered in the box; this suggests feeding at night. Many vacated leaf folds were examined over a wide area on 22nd June suggesting that an earlier search might be more productive. As most larval spinnings were widely spaced, egg-laying may be a travelling chore in contrast to such Agonopterix as heracliana (L.), subpropinquella (Stainton) and alstroemeriana (Clerck) where a number of larvae may be found on one plant.

The moths emerged on 21st and 23rd July, and using a bee smoker on 8th August when the leaves on Sanicula were hidden by a strong growth of Dog's Mercury (Mercurialis) and Enchanter's Nightshade (Circaea), two faded moths were disturbed; a search with a hand lamp in another wood a week later was unsuccessful. Clearly this species has been long established in North Wales and all records from VC 49/50 and the single record from Yorkshire have come from mixed woods on limestone, for Sanicula europaea is mainly a calcareous plant. Of the three Tortricid pupae found in similar folds in the leaves, two were Tortrix viridana (L.) from oak and one was Olindia schumacherana (F.) which probably came from adjacent Mercurialis and Circaea; while uncertain of the Olindia, clearly the Tortrix has a labour saving purpose in using a ready made puparium. — H. N. MICHAELIS, 5 Glan y Mor, Glan Conwy, Colwyn Bay

LL28 5TA.

Danaus plexippus L.: The Monarch in Wales and Cumberland in 1981. — A single specimen was seen and caught on 25th September on the West Cumbrian coast a few hundred yards north of Sellafield railway station. This appears to be the first record of the species for Cumberland. The insect, a female, is now in the possession of Mr. R. Savory, of Millom, but was caught by Mr. F. Downton, of Egremont.

On October 4th, my friend Dr. P. I. Clark was on holiday at Penmaenpool, near Dollgellau, Merioneth, and saw a single specimen flying round the garden for several minutes. — D. W. KYDD, "Gilgar-

ran", 6 Yewbarrow Road, Ulverston, Cumbria LA12 9JS.